

	Healthcare Worker (HCW) Scenario Guidelines COVID-19		
Scenario	Conventional Strategy		
	Testing and Isolation	Leave Provisions	
	e status is defined as 2 doses of COVID-19 vaccine plus a boos		
vaccine OR 2 doses of COVID-19 vaccine AND not greater than 5 months from second dose.			
	al exposures and outbreak testing guidelines, please refer to sc		
Previously COVID-19	NO POCT or PCR required for 90 days post positive PCR test, except for #3 & #4		
Positive Health Care Worker	Note: Testing for scenario #3 & #4 only if greater than 30 days since positive PCR		
	result AND new onset of symptoms. Notify Employee 1. Continue to work	Not applicable.	
#1: Do you or any family members have	 Continue to work HCW should perform POCT test or Abbott ID Now 	Not applicable.	
ONE symptom?	where available. Abbott ID Now is the preferred method		
	for POCT testing while in the workplace and is available		
	at EH on-site testing clinics		
	3. Self-monitor for further symptoms		
	4. If you have fever, loss of taste or smell, follow		
	scenario #3		
#2: Do you have	1. Continue to work	Not applicable.	
another member of	2. Self-monitor for symptoms and if possible, isolate		
your household with	from symptomaticfamily member		
TWO or more	3. If you develop fever, loss of taste or smell follow #3		
symptoms?			
#3: Do you have ONE	, , , ,	Sick time can be used to	
of the following		ensure continuity of salary for	
symptoms? Fever, loss		these absences.	
of taste or smell.		If non-inpatient HCW,	
	•	Manager may offer work from home if the role	
	, 5	permits.	
#4: Do you have TWO or		Sick time can be used to	
more symptoms?		ensure continuity of salary for	
	If PCR negative, return to work when symptoms		
	improve	lf non-inpatient HCW,	
	 If PCR positive, follow scenario #5 	Manager may offer work from	
	3. Notify manager of results	home if the role permits.	
#5: Do you have a		Sick time can be used to	
positive POCT or PCR?		ensure continuity of salary for	
(Day zero is date of		these absences.	
positive POCT or PCR,	at <u>GNB.ca</u>		
whichever comes first.)		If non-inpatient HCW, Managar may offer work from	
		Manager may offer work from home if the role permits.	
	POCT or PCR, whichever comes first		
	6. Notify manager of results		
	7. Return to work for immunocompromised HCW will		
	require and assessment for severity of compromise by the		
	HCW primary care provider if no Primary Health care		
	provider available the HCW can contact <u>eVisitNB</u> for a		
	medical consult. HCW will report severity level to EHS to		
	determine return to work date.		

#6: Are you a close contact of a POCT positive or a confirmed PCR positive case in your home ?	 Remain out of the workplace for 10 days following last contact with the positive case <u>Work isolate</u> for 10 days after return to work Book a PCR test on day 5 and 10 at <u>GNB.ca</u>. You may return to work while waiting for result of day 10 Abbott ID testing is recommended on day 7. If Abbott ID testing not available obtain POCT from manager and perform testing on day 7. Book a PCR at <u>GNB.ca</u> as early as possible if you develop symptoms If you have a positive POCT or PCR follow #5 	Applicable leave of absence to be determined on a case by case basis. Leave provisions not applicable for work isolation. If non-inpatient HCW, Manager may offer work from home if the role permits.
#7: Are you a close contact of a POCT positive or a confirmed PCR positive case in the community and can isolate from the positive case?	 Remain out of the workplace for 5 days following last contact with the positive case <u>Work isolate</u> for 5 days after return to work Book a PCR test at <u>GNB.ca</u> on day 5 and 10. You may return to work while waiting for results. Abbott ID testing is recommended on day 7. If Abbott ID testing not available obtain POCT from manager and perform testing on day 7. If you have a positive POCT or PCR follow #5 	Applicable leave of absence to be determined on a case by case basis. Leave provisions not applicable for work isolation. If non-inpatient HCW, Manager may offer work from home if the role permits.
Occupational Expo	ure	
#8: HCW notified of an occupational exposure to a COVID-19 positive patient or staff member (from 48 hours prior to onset of symptoms or date of POCT/PCR if asymptomatic) Note: not applicable on outbreak units, see scenario #9	 Book a PCR test the day you are notified as well as day 5 and 10 after your initial swab at <u>GNB.ca</u> OR at EH onsite drop in testing clinic. Self-monitor for symptoms If you develop symptoms, do not report to work, book a PCR test at <u>GNB.ca</u> immediately and notify your manager. If PCR positive, refer to scenario #5 If PCR negative, return to work when symptoms improve or refer to scenario #3 or #4. 	Not applicable.
#9: HCW notified of a COVID-19 outbreak and has worked greater than 15 minutes in the outbreak department/unit in the prior 5 days.	 Book a PCR test the day you are notified as well as day 5 and 10 after your initial swab at <u>GNB.ca</u> OR at EH onsite drop in testing clinic. Further testing may be requested per IPC/Infectious Disease Self-monitor for symptoms If you develop symptoms, do not report to work, book a PCR test at <u>GNB.ca</u> immediately and notify your manager. If PCR positive, refer to scenario # 5 If PCR negative, return to work when symptoms improve or refer to scenario #3 or #4. 	Not applicable.